

UNIT – IV Complex integration**Basic concepts:****Connected region:**

A region R is said to be connected when two points of it are connected by a curve which entirely lies inside the region.

Simply connected:

Any closed curve which lies in the region R, can be shrink to a point without learning R is called simply connected region.

Multiply connected region:

A region which is not simply connected is called multiply connected.

Cauchy's integral (Fundamental) theorem:**Statement:**

If $f(z)$ is analytic at every point of the region R bounded by a closed curve C and if $f(z)$ is continuous at all points inside and on closed curve C, then $\int_C f(z) dz = 0$.

Cauchy's integral formula:

If $f(z)$ is analytic inside and on a closed curve C of a simply connected region R and if

a is any point within C, then $f(a) = \frac{1}{2\pi i} \int_C \frac{f(z)}{z-a} dz$.

Cauchy's integral formula for higher derivative:

$$f^n(a) = \frac{ni}{2\pi i} \int_C \frac{f(z)}{(z-a)^{n+1}} dz$$

Taylor's series

A function $f(z)$, analytic inside circle c with centre at $z = a$, can be expanded in the

form $f(z) = f(a) + (z-a) \frac{f'(a)}{1!} + \frac{(z-a)^2}{2!} f''(a) + \dots$ which is convergent at every point inside c.

Laurent's series

Let C_1, C_2 be two concentric circles $|z-a| = R_1$ and $|z-a| = R_2$ where $R_1 < R_2$. Let $f(z)$ be analytic on C_1 and C_2 and in the annular region R between them, then for any point z

in R $f(z) = \sum_{n=0}^{\infty} a_n (z-a)^n + \sum_{n=0}^{\infty} b_n (z-a)^{-n}$ where

$$a_n = \frac{1}{2\pi i} \int_C \frac{f(z)}{(z-a)^{1+n}} dz \text{ and } b_n = \frac{1}{2\pi i} \int_C \frac{f(z)}{(z-a)^{1-n}} dz$$

Standard expansions:

$$1) \quad (1+x)^{-1} = 1 - x + x^2 - x^3 + \dots = \sum_{n=0}^{\infty} (-1)^n x^n \quad \text{if } |x| < 1$$

$$2) \quad (1-x)^{-1} = 1 + x + x^2 + x^3 + \dots = \sum_{n=0}^{\infty} x^n \quad \text{if } |x| < 1$$

$$3) \quad (1+x)^{-2} = 1 - 2x + 3x^2 - 4x^3 + \dots = \sum_{n=0}^{\infty} (-1)^n (n+1)x^n \quad \text{if } |x| < 1$$

$$4) \quad (1-x)^{-2} = 1 + 2x + 3x^2 + 4x^3 + \dots = \sum_{n=0}^{\infty} (n+1)x^n \quad \text{if } |x| < 1$$

$$5) \quad e^x = 1 + x + \frac{x^2}{2!} + \frac{x^3}{3!} + \dots = \sum_{n=0}^{\infty} \frac{x^n}{n!}$$

$$6) \quad \sin x = x - \frac{x^3}{3!} + \frac{x^5}{5!} - \dots$$

$$7) \quad \cos x = 1 - \frac{x^2}{2!} + \frac{x^4}{4!} - \dots$$

Evaluation of Residues

Case(i): if $z = a$ is a simple pole (pole of order 1) the residue of $f(z)$ at $z = a$ is given

$$\text{By } \lim_{z \rightarrow a} (z - a)f(z)$$

Case(ii): if $z = a$ is a pole of order m then Residue of $f(z)$ at $z = a$ is given by

$$\frac{1}{(m-1)!} \lim_{z \rightarrow a} \frac{d^{m-1}}{dz^{m-1}} [(z-a)^m f(z)]$$

Case(iii): if $f(z) = \frac{P(x)}{Q(x)}$ has a simple pole at $z = a$ then residue at $z = a$ is given

$$\text{By } \frac{P(a)}{Q'(a)} \text{ provide } Q'(a) \neq 0.$$

Res $f(z)$ at $z = a =$ coefficient of $\frac{1}{z-a}$ in Laurent series.

Cauchy's Residue theorem:

Statement:

If $f(z)$ is analytic within and on a simple closed curve C except for a finite number of singularities z_1, z_2, \dots, z_n lies inside C and if R_1, R_2, \dots, R_n are the residues of $f(z)$ at these singularities respectively, then

$$\int_C f(z) dz = 2\pi i [R_1 + R_2 + \dots + R_n]$$

Contour Integration:

Type I: $\int_0^{2\pi} F(\sin \theta, \cos \theta) d\theta \rightarrow (1)$

Put $z = e^{i\theta} \Rightarrow \cos \theta = \left(\frac{z^2 + 1}{2z} \right)$ and $\sin \theta = \left(\frac{z^2 - 1}{2iz} \right)$ and C is the unit circle

$|z| = 1, d\theta = \frac{dz}{iz}$ reduce type (1) to the form $\int_C f(z) dz$ which can be evaluated by

Cauchy's residue theorem.

Type II: $\int_{-\infty}^{\infty} \frac{P(x)}{Q(x)} dx$

Consider $\int_C F(z) dz$ where $F(z) = \frac{P(z)}{Q(z)}$ and C is the closed contour consisting of the line segment

$(-R, R)$ together with upper semi circle $|z| = R$

Cauchy's lemma:

If $f(z)$ is continuous function such that $|zf(z)| \rightarrow 0$ uniformly as $|z| = R \rightarrow \infty$ then $\int_C f(z) dz \rightarrow 0$.

Type III: $\int_{-\infty}^{\infty} f(x) \sin mx dx$ (or) $\int_{-\infty}^{\infty} f(x) \cos mx dx$

Jordan's lemma:

If $f(z)$ is continuous function such that $|f(z)| \rightarrow 0$ is uniformly as $|z| = R \rightarrow \infty$ then $\int_C e^{imz} f(z) dz \rightarrow 0$

Evaluate the following by using Cauchy's Integral Theorem

1) $\int_C \frac{z^2 - z + 1}{z - 1} dz$ where C is $|z| = \frac{1}{2}$

Answer : 0

2) $\int_C \frac{z^2 + 1}{z^2 - 1} dz$ where C is $|z - 1| = 1$

Answer : $2\pi i$

3) $\int_C \frac{4 - 3z}{z(z - 1)(z - 2)} dz$ where C is $|z| = \frac{3}{2}$

Answer : $2\pi i$

4) $\int_C \frac{e^{2z}}{(z + 1)^4} dz$ where C is $|z| = 2$

Answer : $\frac{8\pi i}{3e^2}$

5) $\int_C \frac{z}{(z - 1)^2(z + 2)} dz$ where C is $|z| = \frac{3}{2}$

Answer : $\frac{4\pi i}{7}$

6) $\int_C \frac{\sin \pi z^2 + \cos \pi z^2}{(z - 1)(z - 2)} dz$ where C is $|z| = 3$

Answer : $4\pi i$

7) $\int_C \frac{dz}{(z^2 + 4)}$ where C is $|z - i| = 2$

Answer : $\frac{\pi}{16}$

8) $\int_C \frac{z + 1}{z^2 + 2z + 4} dz$ where C is $|z + 1 + i| = 2$

Answer : πi

- 9) If $f(a) = \int_c \frac{4z^2 + z + 5}{(z-a)} dz$ where C is the ellipse $9x^2 + 4y^2 = 36$. Find the values of

$f(1), f(i), f'(-1)$ and $f''(-i)$.

Answer : $20\pi i, 2\pi i(1+i), 14\pi i$ and $16\pi i$

- 10) If $f(a) = \int_c \frac{3z^2 + 7z + 1}{z-a} dz$ where c is $|z| = 2$. Find

$f(3), f(i-1), f'(1-i)$ and $f''(1-i)$

Answer : $0, 2\pi i(8-13i), 2\pi i(13-6i), 12\pi i$

- 11) Evaluate $\int_c \frac{e^z}{z(1-z)^3} dz$ if

(i) 0 lies inside c and 1 lies outside c

(ii) 1 lies inside c and 0 lies outside c

(iii) both 0 and 1 lies inside c.

Problems on Taylor's and Laurent's Series

- 1) Expand $f(z) = \sin z$ as Taylor's series about $z = \frac{\pi}{4}$
- 2) Expand $f(z) = \frac{e^{2z}}{(z-1)^3}$ about $z=1$ using Taylor's series.
- 3) Expand $f(z) = \frac{z^2 - 1}{(z+2)(z+3)}$ in Laurent's series
if (i) $|z| < 2$ (ii) $2 < |z| < 3$ (iii) $|z| > 3$
- 4) Find the Laurent's series of $f(z) = \frac{7z-2}{z(z-2)(z+1)}$ valid in the region
(i) $1 < |z+1| < 3$ (ii) $|z+1| > 3$
- 5) Find the Laurent's series of $f(z) = \frac{z}{(z^2+1)(z^2+4)}$ valid in the region
(i) $1 < |z| < 2$ (ii) $|z| > 2$

Problems on Cauchy's Residue Theorem

1) Determine Residue of the following at each of its poles

a. (i) $f(z) = \cot z$

b. ii) $f(z) = \tan z$

Answer : 1, -1

2) Find the Residue of $f(z) = z^2 \sin\left(\frac{1}{z}\right)$ at $z = 0$

3) Using Cauchy's Residue theorem, evaluate $\int_C \frac{3z^2 + 2}{(z-1)(z^2 + 9)} dz$ Where C is $|z - 2| = 2$

Answer : πi

4) Evaluate $\int_C \frac{dz}{\sin z}$ where C is $|z| = 4$ using Cauchy's residue theorem.

Answer : $2\pi i$

5) Evaluate $\int_C \frac{z \sec z}{(1 - z^2)} dz$ where C is the ellipse $4x^2 + 9y^2 = 9$

Answer : $-2\pi i \sec 1$

6) Evaluate $\int_C \frac{z^2}{z^4 - 1} dz$ where C is $|z| = 2$

Answer : 0

Problems on Contour Integration

(1) $\int_0^{2\pi} \frac{d\theta}{a + b \cos \theta}$ where $a > b > 0$

Answer : $\frac{2\pi}{\sqrt{a^2 - b^2}}$

2) $\int_0^{2\pi} \frac{d\theta}{1 - 2a \sin \theta + a^2}$ $0 < a < 1$

Answer : $\frac{2\pi}{1 - a^2}$

3) Prove that $\int_{-\infty}^{\infty} \frac{x^2 - x + 2}{x^4 + 10x^2 + 9} dx$

Answer : $\frac{5\pi}{12}$

4) Evaluate $\int_{-\infty}^{\infty} \frac{x^2}{(x^2 + a^2)(x^2 + b^2)} dx$

Answer : $\frac{\pi}{a+b}$

5) Prove that $\int_0^{\infty} \frac{\sin mx}{x} dx = \frac{\pi}{2}$, $m > 0$

6) Prove that $\int_0^{\infty} \frac{x \sin x}{x^2 + a^2} dx$

Answer : $\frac{\pi e^{-a}}{2}$